## Nye legemidler som ikke har markedsføringstillatelse

For legemidler oppført på listen gjelder retningslinjens vilkår <u>uavhengig av indikasjon for bruken</u>.

Oppdatert: 29.01.2025

Oppdatert: 29.01.2025	_		
		Orphan	
		medicinal	
		product (Se	
		informasjon	
		om COMP	
		(Komiteen	
		for	
		legemidler	
		mot sjeldne	
		sykdommer)	
		på	
	Indikasjon per nå.	www.legemi	
	(Samt mulig andre indikasjoner i	ddelverket.n	Oppført på
Virkestoff	fremtiden.)	О	listen
abicipar pegol	Treatment of neovascular (wet)		
	age related macular degeneration		
	(AMD)		aug.19
acalabrutinib	Treatment of mantel cell lymphoma -		
	Kun denne indikasjonen		mar.18
	Treatment of wild-type or variant		11101.10
	transthyretin amyloidosis in adult		
acoramidis		х	
acorannuis	patients with cardiomyopathy (ATTR-		
	CM)		mar.24
	Treatment of symptomatic obstructive		
	hypertrophic cardiomyopathy (oHCM) in		
aficamten	adult patients		jan.25
allogeneic bone marrow derived	Treatment of graft-versus-host disease.		
mesenchymal stromal cells, ex-			
vivo expanded		Х	
·			sep.23
alpelisib	Treatment of patients with severe		·
	manifestations of PIK3CArelated		
	overgrowth spectrum - Kun denne	Х	
	indikasjonen		aug.22
anadamtasa alfa	Treatment of congenital thrombotic		aug.22
apadamtase alfa	thrombocytopenic purpura (cTTP) due to		
	, , , , , , , ,	Х	
	ADAMTS13 deficiency		jun.23
arimoclomol	Treatment of Niemann-Pick disease type	x	
	C (NPC)	, and a second s	des.20
artesunate (*Kun styrken 60 mg)	Treatment of severe malaria		
		х	
			feb.21
•		-	- '

des.22
des.22
ues.zz
sep.20
jan.24
okt.20
jan.25
okt.21
/des.21
aug.24
448.21
mar.24
11101.24
-1+ 24/
okt. 24/
nov. 22
des.23
/jun.23
jan.24
jan.24
jan.24
(

attacks of		
		11101.24
		mai.24
To slow disease progression in adult		
Treatment of ATTK amyloidosis	Х	mar.24
		feb.19
		jun.24
·		_
		nov.18
dystrop. Treatment of Duchenne muscular dystrophy	Х	jul.24/ aug.23
Treatment of ambulatory patients aged 3 to 7 years old with Duchenne muscular		,u13
Treatment of neurodegeneration with brain iron accumulation - Kun denne indikasjonen	х	jun.19
advanced or metastatic non squamous nonsmall cell lung cancer (NSCLC)		apr.24
Treatment of anemia assiciated with chronic kidney disease (CKD) in adults		apr.22
marginal zone lymphoma	^	jul.22
Treatment of non-hodgkin lymphoma  Treatment of adult patients with	· ·	mar.18
. , , , ,		160.23
- haemophilia A (congenital factor VIII deficiency) with FVIII inhibitors ≥ 12 years of age haemophilia B (congenital factor IX deficiency) with FIX inhibitors of any age.		feb.23
Indicated for routine prophylaxis to prevent or reduce the frequency of bleeding in patients with		
Indicated for the treatment of malignant ascites		mai.24
dystrophy		0
		jan.25 okt.24
	Indicated for the treatment of malignant ascites  Indicated for routine prophylaxis to prevent or reduce the frequency of bleeding in patients with - haemophilia A (congenital factor VIII deficiency) with FVIII inhibitors ≥ 12 years of age haemophilia B (congenital factor IX deficiency) with FIX inhibitors of any age.  Treatment of non-hodgkin lymphoma  Treatment of adult patients with marginal zone lymphoma  Treatment of adult patients with locally advanced or metastatic non squamous nonsmall cell lung cancer (NSCLC)  Treatment of neurodegeneration with brain iron accumulation - Kun denne indikasjonen  Treatment of ambulatory patients aged 3 to 7 years old with Duchenne muscular dystrop. Treatment of Duchenne muscular dystrop. Treatment of Duchenne muscular dystrophy  Treatment of glioblastoma (GBM)  Indicated for the treatment of cystic fibrosis  Treatment of ATTR amyloidosis  To slow disease progression in adult patients with Alzheimer's disease (AD)  For routine prevention of recurrent	Treatment of duchennes muscular dystrophy  Indicated for the treatment of malignant ascites  Indicated for routine prophylaxis to prevent or reduce the frequency of bleeding in patients with - haemophilia A (congenital factor VIII deficiency) with FVIII inhibitors ≥ 12 years of age haemophilia B (congenital factor IX deficiency) with FIX inhibitors of any age.  Treatment of non-hodgkin lymphoma  Treatment of adult patients with marginal zone lymphoma  Treatment of adult patients with locally advanced or metastatic non squamous nonsmall cell lung cancer (NSCLC)  Treatment of neurodegeneration with brain iron accumulation - Kun denne indikasjonen  Treatment of ambulatory patients aged 3 to 7 years old with Duchenne muscular dystrop. Treatment of Duchenne muscular dystrop. Treatment of Duchenne muscular dystrophy  Treatment of glioblastoma (GBM)  Indicated for the treatment of cystic fibrosis  Treatment of ATTR amyloidosis x  To slow disease progression in adult patients with Alzheimer's disease (AD)  For routine prevention of recurrent attacks of

doxecitine / doxribtimine -	Indicated for the treatment of paediatric		
	and		
	adult patients with thymidine kinase 2		
	deficiency (TK2d) with an age of		
	symptom onset on or before 12 years		
	,,,,,,,		des.24
duvelisib	Treatment of adult patients with small		
	lymphocytic lymphoma (SLL) - <b>Kun</b>	x	
	denne indikasjonen		feb.20
edaravone	Amyotrophic lateral sclerosis (ALS)		jan.18
eflornithine / sulindac	Treatment of familial adenomatous	x	
	polyposis	^	jul.20
	Treatment of high-risk neuroblastoma		
	responsive to prior multiagent,	V	
eflornithine	multimodality therapy - <b>Kun denne</b>	Х	
	indikasjonen		jan.25
	for the treatment of moderate to severe		
elinzanetant	vasomotor symptoms (VMS)		des.24
enasidenib			
	Treatment of acute myeloid leukaemia		mar.18
entolimod			nov.17
	Treatment of acute radiation syndrome	Х	
eplontersen	Treatment of adult patients with		
•	polyneuropathy associated with		
	hereditary transthyretin-mediated	Х	
	amyloidosis (ATTRv)		nov.23
eteplirsen	Treatment of duchennes muscular		okt.24
'	dystrophy		
ferric citrate coordination	Treatment of iron deficiency anaemia in		
complex	adult chronic kidney disease (CKD)		
·	patients with elevated serum		
	phosphorus levels		jun.24
ferumoxytol	Intravenous treatment of iron deficiency		-
,	anaemia (IDA)		jun.22
garadacimab	Routine prevention of attacks of		•
	hereditary angioedema (HAE)	Х	des.23
glutamine	Treatment of sickle cell disease		mar.18
golodirsen	Treatment of duchennes muscular		okt.24
80.00	dystrophy		
govorestat	Treatment of adults and		
801010300	children aged 2 years and older with a		
	confirmed diagnosis of classic	х	
	galactosemia		jan.24
idebenone	Treatment of respiratory dysfunction in		۱ ۵۰۰۰۰ ۱
laebenone	patients with Duchenne muscular		
	dystrophy (DMD) not using		
	glucocorticoids - <b>Kun denne</b>	x	
	indikasjonen		
			aug.19
			446.13

transfusion-dependent anaemia due to very low, low and intermediate-risk myelodysplastic syndromes (MDS) with ring sideroblasts  imlunestrant  Monotherapy and in Combination with Verzenio® (abemaciclib) in Patients with ER+, HER2- Advanced Breast Cancer  inavolisib  Treatment of adult patients with PIK3CA-mutated, hormone receptor (HR3-positive, human epidermal growth factor receptor 2 (HER2)-negative, locally advanced or metastatic breast cancer  infigratinib  Treatment of cholangiocarcinoma  X  ipatasertib  Treatment of breastcancer indicated in adults and children from birth for chronic treatment of Niemann-Pick Type C (NPC)  lazertinib  Treatment of adult patients with advanced nonsmall cell lung cancer (NSCLC)  lecanemab  Indicated as a disease modifying treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  Ienadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  Ieniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  Treatment of unresectable or metastatic melanoma  Iinvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor-Induced Acneiform Rash			Treatment of adult patients with	imetelstat
imlunestrant  Monotherapy and in Combination with Verzenio® (abemaciclib) in Patients with ER+, HER2- Advanced Breast Cancer  linavolisib  Treatment of adult patients with PIK3CA-mutated, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, locally advanced or metastatic breast cancer  infigratinib  Treatment of cholangiocarcinoma  x  ipatasertib  Treatment of breastcancer  H-acetylleucine  indicated in adults and children from birth for chronic treatment of Niemann-Pick Type C (NPC)  lazertinib  Treatment of adult patients with advanced onsmall cell lung cancer (NSCLC)  lecanemab  Indicated as a disease modifying treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  lenadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  leniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  Iffileucel  Treatment of unresectable or metastatic melanoma  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor-Induced Acneiform Rash			transfusion-dependent anaemia due to	
imlunestrant  Monotherapy and in Combination with Verzenio* (abemaciclib) in Patients with ER+, HER2- Advanced Breast Cancer  inavolisib  Treatment of adult patients with PIK3CA- mutated, hormone receptor (HR)- positive, human epidermal growth factor receptor 2 (HER2)-negative, locally advanced or metastatic breast cancer  infigratinib  Treatment of cholangiocarcinoma  x  Treatment of breastcancer I-acetylleucine  indicated in adults and children from birth for chronic treatment of Niemann- Pick Type C (NPC)  lazertinib  Treatment of adult patients with advanced nonsmall cell lung cancer (NSCLC)  lecanemab  Indicated as a disease modifying treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  lenadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  leniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  Iffileucel  Treatment of unresectable or metastatic melanoma  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash		v	very low, low and intermediate-risk	
imlunestrant  Monotherapy and in Combination with Verzenio® (abemaciclib) in Patients with ER+, HER2- Advanced Breast Cancer  inavolisib  Treatment of adult patients with PIK3CA-mutated, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, locally advanced or metastatic breast cancer  infigratinib  Treatment of cholangiocarcinoma  x  ipatasertib  Treatment of breastcancer I-acetylleucine  indicated in adults and children from birth for chronic treatment of Niemann-Pick Type C (NPC)  lazertinib  Treatment of adult patients with advanced nonsmall cell lung cancer (NSCLC)  lecanemab  Indicated as a disease modifying treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  lenadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  leniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  Iifileucel  Treatment of unresectable or metastatic melanoma  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash		Х	myelodysplastic syndromes (MDS) with	
Verzenio® (abemaciclib) in Patients with ER+, HER2- Advanced Breast Cancer  inavolisib  Treatment of adult patients with PIK3CA-mutated, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, locally advanced or metastatic breast cancer  infigratinib  Treatment of cholangiocarcinoma  x  ipatasertib  I-acetylleucine  indicated in adults and children from birth for chronic treatment of Niemann-Pick Type C (NPC)  lazertinib  Treatment of adult patients with advanced nonsmall cell lung cancer (NSCLC)  lecanemab  Indicated as a disease modifying treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  lenadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  leniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  Treatment of unresectable or metastatic melanoma  linvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash			ring sideroblasts	
Verzenio® (abemaciclib) in Patients with ER+, HER2- Advanced Breast Cancer  inavolisib  Treatment of adult patients with PIK3CA-mutated, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, locally advanced or metastatic breast cancer  infigratinib  Treatment of cholangiocarcinoma  x  ipatasertib  I-acetylleucine  indicated in adults and children from birth for chronic treatment of Niemann-Pick Type C (NPC)  lazertinib  Treatment of adult patients with advanced nonsmall cell lung cancer (NSCLC)  lecanemab  Indicated as a disease modifying treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  lenadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  leniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  Treatment of unresectable or metastatic melanoma  linvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash	nov.23			
ER+, HER2- Advanced Breast Cancer  inavolisib  Treatment of adult patients with PIK3CA- mutated, hormone receptor (HR)- positive, human epidermal growth factor receptor 2 (HER2)-negative, locally advanced or metastatic breast cancer  infigratinib  Treatment of cholangiocarcinoma  x  ipatasertib  Treatment of breastcancer Inacetylleucine  indicated in adults and children from birth for chronic treatment of Niemann- Pick Type C (NPC)  lazertinib  Treatment of adult patients with advanced nonsmall cell lung cancer (NSCLC)  lecanemab  Indicated as a disease modifying treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  lenadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  leniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  Treatment of unresectable or metastatic melanoma  linvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash			• •	imlunestrant
inavolisib  Treatment of adult patients with PIK3CA- mutated, hormone receptor (HR)- positive, human epidermal growth factor receptor 2 (HER2)-negative, locally advanced or metastatic breast cancer  infigratinib  Treatment of cholangiocarcinoma  x  ipatasertib  Treatment of breastcancer I-acetylleucine  indicated in adults and children from birth for chronic treatment of Niemann- pick Type C (NPC)  Iazertinib  Treatment of adult patients with advanced nonsmall cell lung cancer (NSCLC)  Iecanemab  Indicated as a disease modifying treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  Ienadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  Ieniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  Iifileucel  Treatment of urresectable or metastatic melanoma  Iinvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash				
mutated, hormone receptor (HR)- positive, human epidermal growth factor receptor 2 (HER2)-negative, locally advanced or metastatic breast cancer  infigratinib  Treatment of cholangiocarcinoma  x  ipatasertib  Treatment of breastcancer  l-acetylleucine  indicated in adults and children from birth for chronic treatment of Niemann- Pick Type C (NPC)  lazertinib  Treatment of adult patients with advanced nonsmall cell lung cancer (NSCLC)  lecanemab  Indicated as a disease modifying treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  lenadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  leniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  lifileucel  Treatment of urresectable or metastatic melanoma  linvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash	jan.25		ER+, HER2- Advanced Breast Cancer	
positive, human epidermal growth factor receptor 2 (HER2)-negative, locally advanced or metastatic breast cancer infigratinib  Treatment of cholangiocarcinoma  x  ipatasertib  Treatment of breastcancer I-acetylleucine indicated in adults and children from birth for chronic treatment of Niemann-Pick Type C (NPC)  Iazertinib  Treatment of adult patients with advanced nonsmall cell lung cancer (NSCLC)  Iecanemab  Indicated as a disease modifying treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  Ienadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  Ieniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  Iifileucel  Treatment of unresectable or metastatic melanoma  Iinvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash	Januar		Treatment of adult patients with PIK3CA-	inavolisib
factor receptor 2 (HER2)-negative, locally advanced or metastatic breast cancer  infigratinib  Treatment of cholangiocarcinoma  x  ipatasertib  Treatment of breastcancer indicated in adults and children from birth for chronic treatment of Niemann-Pick Type C (NPC)  lazertinib  Treatment of adult patients with advanced nonsmall cell lung cancer (NSCLC)  lecanemab  Indicated as a disease modifying treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  lenadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  leniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  lifileucel  Treatment of unresectable or metastatic melanoma  linvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash			mutated, hormone receptor (HR)-	
locally advanced or metastatic breast cancer			positive, human epidermal growth	
locally advanced or metastatic breast cancer			factor receptor 2 (HER2)-negative,	
cancer infigratinib  Treatment of cholangiocarcinoma  X  ipatasertib  Treatment of breastcancer indicated in adults and children from birth for chronic treatment of Niemann-Pick Type C (NPC)  lazertinib  Treatment of adult patients with advanced nonsmall cell lung cancer (NSCLC)  lecanemab  Indicated as a disease modifying treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  lenadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  leniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  lifileucel  Treatment of unresectable or metastatic melanoma  linvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash				
ipatasertib  I-acetylleucine  I-acetylleucine  Indicated in adults and children from birth for chronic treatment of Niemann-Pick Type C (NPC)  Iazertinib  Treatment of adult patients with advanced nonsmall cell lung cancer (NSCLC)  Iecanemab  Indicated as a disease modifying treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  Ienadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  Ieniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  Iifileucel  Treatment of unresectable or metastatic melanoma  Iinvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash	jun.24		-	
ipatasertib  Treatment of breastcancer  I-acetylleucine  indicated in adults and children from birth for chronic treatment of Niemann-Pick Type C (NPC)  Iazertinib  Treatment of adult patients with advanced nonsmall cell lung cancer (NSCLC)  Iecanemab  Indicated as a disease modifying treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  Ienadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  Ieniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  Iifileucel  Treatment of unresectable or metastatic melanoma  Iinvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash		Х	Treatment of cholangiocarcinoma	infigratinib
I-acetylleucine indicated in adults and children from birth for chronic treatment of Niemann-Pick Type C (NPC)  Iazertinib Treatment of adult patients with advanced nonsmall cell lung cancer (NSCLC)  Iecanemab Indicated as a disease modifying treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  Ienadogene nolparvovec Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  Ieniolisib Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  Iifileucel Treatment of unresectable or metastatic melanoma  Iinvoseltamab Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014 Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash	des.21		To a to a set of the control of the	
birth for chronic treatment of Niemann- Pick Type C (NPC)  Izertinib  Treatment of adult patients with advanced nonsmall cell lung cancer (NSCLC)  Iecanemab  Indicated as a disease modifying treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  Ienadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  Ieniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  Iifileucel  Treatment of unresectable or metastatic melanoma  Iinvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash	mar.18			'
Pick Type C (NPC)				l-acetylleucine
lazertinib  Treatment of adult patients with advanced nonsmall cell lung cancer (NSCLC)  Indicated as a disease modifying treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  Ienadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  Ieniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  Iifileucel  Treatment of unresectable or metastatic melanoma  Iinvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash				
advanced nonsmall cell lung cancer (NSCLC)  lecanemab  Indicated as a disease modifying treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  lenadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  leniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  lifileucel  Treatment of unresectable or metastatic melanoma  linvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash	jul.24			
Indicated as a disease modifying treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)    Ienadogene nolparvovec				lazertinib
Indicated as a disease modifying treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  Ienadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  Ieniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  Iifileucel  Treatment of unresectable or metastatic melanoma  Iinvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash				
treatment in adult patients with Mild Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  lenadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  leniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  lifileucel  Treatment of unresectable or metastatic melanoma  linvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash	mar.24		(NSCLC)	
Cognitive Impairment due to Alzheimer's disease and Mild Alzheimer's disease (Early Alzheimer's disease)  Ienadogene nolparvovec Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  Ieniolisib Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  Iifileucel Treatment of unresectable or metastatic melanoma  Iinvoseltamab Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014 Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash			Indicated as a disease modifying	lecanemab
disease and Mild Alzheimer's disease (Early Alzheimer's disease)  lenadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  leniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  lifileucel  Treatment of unresectable or metastatic melanoma  linvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash			treatment in adult patients with Mild	
Alzheimer's disease (Early Alzheimer's disease)  lenadogene nolparvovec  Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  leniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  lifileucel  Treatment of unresectable or metastatic melanoma  linvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash			Cognitive Impairment due to Alzheimer's	
lenadogene nolparvovec Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  leniolisib Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  lifileucel Treatment of unresectable or metastatic melanoma  linvoseltamab Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014 Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash			disease and Mild	
lenadogene nolparvovec Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  Ieniolisib Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  Iifileucel Treatment of unresectable or metastatic melanoma  Iinvoseltamab Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014 Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash			Alzheimer's disease (Early Alzheimer's	
Treatment of vision loss due to Leber Hereditary Optic Neuropathy (LHON)  Ieniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  Treatment of unresectable or metastatic melanoma  Iinvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash	feb.23		disease)	
Hereditary Optic Neuropathy (LHON)  leniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  lifileucel  Treatment of unresectable or metastatic melanoma  linvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash			_	lenadogene nolparvovec
leniolisib  Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  lifileucel  Treatment of unresectable or metastatic melanoma  linvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash		х		
Treatment of activated phosphoinositide 3-kinase delta syndrome (APDS)  lifileucel Treatment of unresectable or metastatic melanoma  linvoseltamab Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014 Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash	nov.20		Hereditary Optic Neuropathy (LHON)	
3-kinase delta syndrome (APDS)  lifileucel Treatment of unresectable or metastatic melanoma  linvoseltamab Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014 Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash			T	leniolisib
lifileucel  Treatment of unresectable or metastatic melanoma  linvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash		х		
melanoma  linvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash	nov.22		, , , ,	
linvoseltamab  Monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash				lifileucel
patients with relapsed or refractory multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash	sep.24			
multiple myeloma  LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash			·	linvoseltamab
LUT014  Topical BRAF Inhibitor Therapy for EGFR Inhibitor–Induced Acneiform Rash			·	
Topical BRAF Inhibitor Therapy for EGFR Inhibitor—Induced Acneiform Rash	mar.24		multiple myeloma	
Inhibitor–Induced Acneiform Rash			Tonical DDAE Inhibitor Thereas for ECED	LU1014
	22			
LEGGERGORE OF MULLINATIONS	nov.23			
mavorixafor Treatment of WHIM syndrome x	jan.25	Х	· ·	
mirdametinib Treatment of neurofibromatosis type 1	sep.24		rreatment of neuronbromatosis type 1	ının udinletifild

mobocertinib		Ī	
	Treatment of adult patients with		
	epidermal growth factor receptor (EGFR)		
	exon 20 insertion mutation-positive		
	locally advanced or metastatic non-small		
	cell lung cancer (NSCLC).		jan.21
mozafancogene autotemcel	Treatment of paediatric patients with	Х	
	Fanconi Anaemia Type A	^	mai.24
nadofaragene firadenovec			
	Treatment of adult patients with high-		
	grade		
	(HG), Bacillus Calmette-Guérin (BCG)-		
	unresponsive non-muscle invasive		
	bladder		
	cancer (NMIBC).,		jan.25
naporafenib	Treatment of advanced solid tumours		okt.21
n a d a siva n	including ovarian cancer		OKL.21
nedosiran	Til behandling av primær hyperoksaluri.		mai.24
nipocalimab	treatment of generalised Myasthenia		
	Gravis		okt.24
nemolizumab	Treatment of moderate-to-severe atopic		
	dermatitis and for the treatment of		
	prurigo nodularis		mar.24
nirogacestat	Treatment of desmoid tumours	Х	apr.24
nogapendekin alfa / Inbakicept			
	Treatment of adult patients with BCG-		
	unresponsive non-muscle invasive		
	bladdercancer (NMIBC) with carcinoma		
	in situ (CIS)		
	with or without papillary tumours		jan.25
obecabtagene autoleucel	Treatment of patients with relapsed or		
	refractory B cell precursor acute	х	
	lymphoblastic leukaemia (ALL)		mai.24
obeticholic acid			
	Improvement of liver fibrosis and		
	resolution of steatohepatitis in adult		
	patients with significant liver fibrosis due		
	to nonalcoholic steatohepatitis (NASH).		
	Kun denne indikasjonen		aug.20
obnitix	Treatment of RR-aGVHD	<del>                                     </del>	apr.24
odevixibat			
odevixibat	Treatment of cholestatic pruritus in		
odevixibat	Treatment of cholestatic pruritus in Alagille syndrome (ALGS) <b>Kun denne</b>		
odevixibat	·		jan.24
	Alagille syndrome (ALGS) <b>Kun denne</b>		jan.24
odevixibat odronextamab	Alagille syndrome (ALGS) Kun denne indikasjonen		jan.24
	Alagille syndrome (ALGS) <b>Kun denne indikasjonen</b> Treatment of blood cancers (follicular		jan.24
	Alagille syndrome (ALGS) <b>Kun denne indikasjonen</b> Treatment of blood cancers (follicular lymphoma (FL) or diffuse large B cell		jan.24 mai.24

olezarsen	Treatment of familial chylomicronemia		
	, in the second of the second		sep.24
omadacycline tosylate			
	Treatment of community-acquired		
	bacterial pneumonia (CABP) and acute		
	bacterial skin and skin structure		
	infections (ABSSSI) in adults		okt.18
omecamtiv mecarbil	Treatment of adult patients with		
	symptomatic chronic heart failure and		
	reduced ejection fraction less than 30%		
			feb.23
oportuzumab monatox			
	Treatment and prevention of recurrence		
	of carcinoma-in-situ (CIS) of the urinary		
	bladder and prevention of recurrence of		
	high grade Ta and/or T1 papillary		
	tumours		apr.21
parsaclisib	Treatment of adult patients with		
	relapsed or refractory marginal zone	х	
	lymphoma (MZL)		feb.22
pegcetacoplan	Treatment of geographic atrophy (GA)		
	secondary to age-related macular		
	degeneration (AMD) - <b>Kun denne</b>	Х	
	indikasjonen		feb.23
pevonedistat			
	Higher-risk chronic myelomonocytic		
	leukemia (HR-CMML), low-blast acute		
	myeloid leukemia (LB-AML) and higher		
	risk myelodysplastic syndromes		aug.20
plazomicin			
	Treatment of symptomatic tenosynovial		
	giant cell tumor (TGCT), also known as		
	pigmented villonodular synovitis (PVNS)		
	and giant cell tumor of the tendon		
	sheath (GCT-TS), where surgical	Х	
	resection is potentially associated with		
	worsening functional limitation or		
	severe morbidity		
	severe morbidity	1	mar.19
plazomicin	Treatment of Complicated urinary tract		111011.13
P	infection (cUTI), including		
	pyelonephritis; treatment of		
	Bloodstream infection (BSI);		
	treatment of infections due to		
	Enterobacteriaceae		nov.18
pridopidine	Treatment of Huntington's disease		
pridopidirie	ricatificht of Hulltington's disease	X	sep.24

resmetirom			
	Treatment of adults with nonalcoholic steatohepatitis (NASH)/metabolic dysfunctionassociated steatohepatitis		
	(MASH) with liver fibrosis		apr.24
resminostat	Treatment of patients with advanced stage mycosis fungoides (MF) and Sézary syndrome (SS)		apr.24
retifanlimab	Treatment of locally advanced or metastatic squamous carcinoma of the anal canal (SCAC) who have progressed on or who are intolerant of platinum (kun denne indikasjonen)	x	mar.21
rilonacept	treatment of idiopathic pericarditis	Х	okt.24
rilzabrutinib	Treatment of persistent or chronic immune thrombocytopenia (ITP)	X	des.24
romidepsin	Treatment of peripheral T-cell lymphoma (PTCL) in adult patients who have received at least one prior therapy.		feb.22
rovalpituzumab tesirine	Small cell lung cancer (SCLC)		jan.18
savolitinib	monotherapy for the treatment of adult patients with relapsed or refractory multiple myeloma		mar.23
sebetralstat	treatment of hereditary angioedema (HAE) attacks in adult and adolescents aged 12 years and older.	x	sep.24
sela del parly sin di hydrat	A peroxisome proliferator receptor delta (PPARδ) activator indicated for the treatment of primary biliary cholangitis (PBC) including pruritus in adults without cirrhosis or with compensated cirrhosis (Child-Pugh A) in combination with ursodeoxycholic acid (UDCA) who have an inadequate response to UDCA alone, or as monotherapy in those unable to tolerate UDCA.		apr.24
sepiapterin	Treatment of hyperphenylalaninemia		
	(HPA) in adult and paediatric patients with phenylketonuria (PKU) patients with phenylketonuria (PKU).		jun.24

serplulimab	First-line treatment of adult patients		1
	with extensive-stage small cell lung	x	
	cancer (ES-SCLC)		apr.23
siponimod	Indicated for the treatment of		
	thrombocytopenia - <b>Kun denne</b>		
	indikasjonen		okt.18
sodium phenylbutyrate/	Treatment of amyotrophic lateral	х	
ursodoxicoltaurine	sclerosis (ALS)	^	apr.22
surufatinib	Treatment of progressive		
	neuroendocrine tumours		jul.21
tazemetostat			
	Treatment of malignant mesothelioma		
	and Epithelioid sarcoma		mai.21
tegomil fumarat	Treatment of multiple sclerosis		mai.24
teplizumab	To delay both the onset of Stage 3 type		
	1 diabetes (T1D) and the progression of		
	Stage 3 T1D. Utsette		
	klinisk debut av type 1 diabetes		
			jan.25/ jul.24
teprotumumab	Treatment of moderate to severe		
·	Thyroid Eye Disease (TED)		jun.24
tiratricol	Treatment of monocarboxylate		
	transporter 8 (MCT8) deficiency	Х	nov.23
tisotumab vedotin	treatment of adult patients with		
	recurrent or metastatic cervical cancer		
	with disease progression on or after		
	systemic therapy		mar.24
trastuzumabduocarmazine	Treatment of HER2 (Human Epidermal		
	Growth Factor Receptor 2)-positive		
	metastatic breast		aug.22
trofinetide	Treatment of Rett syndrome in adults		Ü
	and paediatric patients	Х	
	2 years of age and older	^	jan.25
troriluzole	, ,		Ja5
	Treatment of adult patients with	х	
	spinocerebellar ataxia genotype 3 (SCA3	Α	nov.23
valemetostat	, , , , , , , , , , , , , , , , , , ,		
varemetostat	Ved tilbakevendende/refraktær (R/R)		
	adult T-cell leukemia/lymphoma (ATLL).		apr.22
veliparib	7 7 15 2 2 3 3 3 3 3 3		Sp.:22
venpario	Treatment of BRCA 1, BRCA 2, BARD1		
	and/orr PALB2 mutated cancer		mar.18
viltolarsen	Treatment of duchennes muscular		okt.24
Viitolai Scii	dystrophy		
vimseltinib	. /		
VIIIISCIUIIIS	Treatment of adult patients with		
	tenosynovial giant cell tumour (TGCT)	х	
	who are not amenable to surgery		aug.24
Vorasidenib	Til behandling av gliom		mai.24
งบาสรเนยเแม	Th behanding av gilotti		IIIdi.24

X