Nye legemidler som ikke har markedsføringstillatelse

For legemidler oppført på listen gjelder retningslinjens vilkår <u>uavhengig av indikasjon for bruken.</u>

Oppdatert: 08.05.2018

Oppdatert: 08.05.2018			
	Indikasjon per nå.	Orphan medicinal product (* Se	
	(Samt mulig andre indikasjoner i fremtiden.)	informasjon nederst)	Oppført på
Virkestoff			listen
Axalimogene filolisbac	Treatment of cervical cancer		mai.18
	Treatment of adults with		
	phenylketonuria (PKU) who have		
	inadequate blood phenylalanine control		
Pegvaliase		х	mai.18
	Monotherapy, indicated for the		
	treatment of patients with metastatic		
	cutaneous squamous cell carcinoma		
Cemiplimab			mai.18
	Treatment and prophylaxis of bleeding in		
	patients with haemophilia A		
Turoctocog alfa pegol		x	mai.18
Glutamine	Treatment of sickle cell disease		mar.18
	First-line treatment of adults with locally		
	advanced or metastatic non-small cell		
	lung cancer (NSCLC) with epidermal		
	growth factor receptor (EGFR)-activating		
Dacomitinib	mutations.,		mar.18
Trientine dihydrochloride	Treatment of Wilson's disease	х	mar.18
	Treatment for long-term enzyme		
	replacement therapy in patients with		
Velmanase alfa	alpha-mannosidosis	х	mar.18
Selumetinib	Neurofibromatosis		mar.18
	Treatment of amyotrophic lateral		
Masitinib	sclerosis (ALS). Se fotnote under**		mar.18
Avapritinib	Treatment of mastocytosis and GIST		mar.18
Ipatasertib	Treatment of breastcancer		mar.18
	Treatment of non- small cell lungcancer		
	(NSCLC) and glioblastoma		
Veliparib			mar.18
'	Treatment of breastcancer and		
Talazoparib	prostatecancer		mar.18
	Treatment of non-hodgkin lymphoma		
Copanlisib			mar.18
	Treatment of acute myeloid leukaemia		
Enasidenib			mar.18
A salah mutik ti	Treatment of mantel cell lymphoma		40
Acalabrutinib	· ·		mar.18
	Conditioning treatment prior to		
	allogeneic haematopoietic stem cell		
	transplantation (alloHSCT)		. =
Treosulfan	transplantation (alloHSCT)	Х	mar.18

	Prevention of major cardiovascular		
Canakinumab	events		mar.18
Lusutrombopag	Treatment of thrombocytopenia		mar.18
Lusutrombopag	Treatment of adult patients with		11101.120
	anaplastic lymphoma kinase (ALK)-		
	positive advanced non-small cell lung		
Lorlatinib	cancer (NSCLC)		mar.18
	Treatment of acute lymphoblastic		
Asparaginase	leukaemia	X	mar.18
	Adjunctive therapy of seizures associated		
	with Lennox-Gastaut syndrome (LGS) or		
	Dravet syndrome (DS),		
Cannabidiol		Х	mar.18
	Prevention of episodic and chronic		
Fremanezumab	migraine		mar.18
	Avatrombopag is indicated for the		
	treatment of thrombocytopenia in adult		
	patients with chronic liver disease who		
	are scheduled to undergo a procedure.		
Avatrompag Maleat			mar.18
	Routine prevention of angioedema		
	attacks and the control of symptoms of		
	hereditary angioedema (HAE) in patients		
Lanadelumab	aged 12 years and older	X	feb.18
	Metastatic Cutaneous Squamous Cell		
	Carcinoma (CSCC) indicated for the		
	treatment of patients with metastatic		
	cutaneous squamous cell carcinoma		
	(mCSCC), or with locally advanced		
	cutaneous squamous cell carcinoma		
Dog 2010	(laCSCC) who are not candidates for		fab 10
Regn2810	Small cell lung cancer (SCLC) -		feb.18
	, ,		
Rovalpituzumab tesirine	maintenance following first-line platinum chemotherapy		jan.18
Edaravone	Amyotrophic lateral sclerosis (ALS)		jan.18
	Treatment of hereditary transthyretin-		janiso
Patisiran	mediated amyloidosis	X	jan.18
T delon di	Induction of response in adult patients		jamze
	with granulomatosis with polyangiitis		
	(Wegener's) (GPA) or microscopic		
Avacopan	polyangiitis (MPA	X	jan.18
Romosozumab	Treatment of osteoporosis	<u> </u>	jan.18
	Intended for the treatment of adult men		, , ,
	with NM-CRPC who are at high risk of		
	developing metastatic disease.		
Apalutamide	' -		jan.18
			,

r			1
	Treatment of adults infected with HIV-1		
	without past or present evidence of viral		
	resistance to treatment of adults infected		
	with HIV-1 without past or present		
	evidence of viral resistance to doravirine,		
Doravirine			des.17
	Treatment of adults infected with HIV-1		
	without past or present evidence of viral		
Doravirine / lamivudine /	resistance to doravirine, lamivudine, or		
tenofovir disoproxil	tenofovir		des.17
Galcanezumab	Prophylaxis of migraine		des.17
Galcariczarriab	Treatment of transthyretin amyloidosis		uc3.17
Inotersen	(hATTR)	v	dos 17
inotersen	·	X	des.17
NA	Diagnosis of Adult growth hormone		.147
Macimorelin	deficiency (AGHD)		des.17
L	To support the endogenous clotting		
Human fibrinogen / human	process and increase of haemostasis in		
thrombin	surgical procedures		des.17
	Treatment of adults with high-risk acute		
Daunorubicin / cytarabine	myeloid leukaemia (AML)	Х	des.17
	Treatment of acute radiation syndrome		
Entolimod		X	nov.17
	Treatment of cutaneous T-cell lymphoma		
Mogamulizumab		X	nov.17
	Treatment of locally advanced,		
	unresectable non-small cell lung cancer		
	(NSCLC).		
Durvalumab	Urotelial cancer (mars 2018)		okt.17
	Treatment and prophylaxis of		0.1101.21
Damoctocog alfa pegol	haemophilia A	X	okt.17
Damostocog and pego.	Treatment of adults infected with human		OKC.17
Bictegravir / emtricitabine /	immunodeficiency virus-1 (HIV-1)		
tenofovir alafenamide	illimationeticiency virus-1 (Titv-1)		jul.17
teriorovii alarenamide	Treatment of disease-related		Jui.17
	splenomegaly and control of symptoms		
	in patients with primary myelofibrosis		
	(PMF), post-polycythemia vera		
	myelofibrosis (PPV-MF), or post-essential		
	thrombocythemia myelofibrosis (PET-		
	MF) who have thrombocytopenia		
	(platelet counts ≤100,000 /μL)		
Pacritinib			jul.17
i acritinio	Routine prophylaxis to prevent bleeding		jui.17
	or reduce the frequency of bleeding		
	episodes in patients with haemophilia A		
	(congenital factor VIII deficiency) with		
Francisia	factor VIII inhibitors.		:. 1 4 =
Emicizumab			jul.17

	Temporary improvement in the		
	appearance of moderate to severe		
Botulinum toxin type a	vertical lines between the eyebrows		jul.17
Meropenem / vaborbactam	Treatment of infections		jul.17
	The treatment or patients with CF aged		
	12 years and older who have at least one		ļ
	F508del mutation in the CFTR gene, and a		
	second mutation that is responsive to		
	tezacaftor/ivacaftor		
Ivacaftor/ tezacaftor			jun.17
	Intended for the treatment of adult		
	patients with diffuse large B-cell		
	lymphoma (DLBCL) who have not		
Autologous T cells transduced	responded to their prior therapy, or have		
with retroviral vector encoding	had disease progression after autologous		
an anti-CD19 CD28/CD3-zeta	stem cell transplant (ASCT)		
chimeric antigen receptor		х	jun.17
	Indicated for prophylaxis of migraine in		
Erenumab	adults		jun.17
Dolutegravir / rilpivirine	Treatment of HIV		jun.17
	Treatment of von Willebrand Disease		
Vonicog alfa	(VWD)	x	jun.17
* Se informasjon om COMP			
(komiteen for legemidler mot			
sjeldne sykdommer) på			
www.legemiddelverket.no			
**Fotnote/ Masitinib:			
http://www.ema.europa.eu/do			
cs/en_GB/document_library/Su			
mmary_of_opinion			
_Initial_authorisation/human/0			
04398/WC500247742.pdf			